

Instructions for Miralax 2-Day Bowel Prep

**Dartmouth Hitchcock Manchester
Endoscopy Center**
100 Hitchcock Way, 3rd Floor
Manchester, NH
(603) 629-8299

Elliot Hospital
1 Elliot Way, Pavilion Entrance- 2nd Floor
Manchester, NH
(603) 663-2170

Notify D-H Gastro Scheduling at 603-629-8299 immediately of any health changes, such as a cold with a fever, within 48 hours of your scheduled procedure, or give us a call if you have any questions.

Patient Name:	Date:	Provider:
If you're scheduled at D-H, you will receive a call 1 week prior to your procedure from our scheduler to assign your arrival time. Since schedules are subject to change, "myD-H" online often times is not a reliable source for an accurate time to arrive— please rely solely on a call from our scheduler for an accurate arrival time.		

<i>7 DAYS PRIOR</i>																																											
<p>PURCHASE:</p> <ul style="list-style-type: none"> • 2 bottles of Miralax or the generic equivalent (238 grams) • 1 bottle of Miralax (119 grams) • 1 box of Dulcolax laxative tabs - 5 mg each <p>TAKE 17 grams (1 capful) of Miralax once daily for a week before you start the prep and follow a low fiber diet.</p> <p>CHECK MEDICATIONS: <u>Do not stop taking all your medications.</u></p> <p>Call your prescribing physician for instructions on how to take your medications prior to procedure if you take any blood thinning medications, including but not limited to:</p> <table> <tr> <td>Aggrenox</td> <td>Coumadin</td> <td>Effient</td> <td>Plavix</td> </tr> <tr> <td>Pradaxa</td> <td>Ticlid</td> <td>Xaralto</td> <td>Lovenox</td> </tr> <tr> <td>Jantoven</td> <td></td> <td></td> <td></td> </tr> </table>	Aggrenox	Coumadin	Effient	Plavix	Pradaxa	Ticlid	Xaralto	Lovenox	Jantoven				<p>Do NOT take:</p> <table> <tr> <td>Advil</td> <td>Alka-seltzer</td> <td>Anaprox</td> <td>Ibuprofen</td> </tr> <tr> <td>Arthrotec</td> <td>Vitamin E</td> <td>Iron pills</td> <td>Mobic</td> </tr> <tr> <td>Escorting</td> <td>Feldene</td> <td>Relafen</td> <td>Multivitamins</td> </tr> <tr> <td>Motrin</td> <td>Narcosis</td> <td>Orudis</td> <td>Unpin</td> </tr> <tr> <td>Aleve</td> <td>Daypro</td> <td>Iodine</td> <td>Naproxen</td> </tr> <tr> <td>Fish oil (Omega -3)</td> <td></td> <td>Lodine</td> <td>Piroxicam</td> </tr> </table> <p>You may continue to take:</p> <table> <tr> <td>Celebrex</td> <td>Maintenance medications</td> </tr> <tr> <td>Methotrexate</td> <td>Plaquenil</td> </tr> <tr> <td>Seizure medications</td> <td>Tylenol</td> </tr> </table> <p>*DO NOT STOP TAKING ASPIRIN</p>	Advil	Alka-seltzer	Anaprox	Ibuprofen	Arthrotec	Vitamin E	Iron pills	Mobic	Escorting	Feldene	Relafen	Multivitamins	Motrin	Narcosis	Orudis	Unpin	Aleve	Daypro	Iodine	Naproxen	Fish oil (Omega -3)		Lodine	Piroxicam	Celebrex	Maintenance medications	Methotrexate	Plaquenil	Seizure medications	Tylenol
Aggrenox	Coumadin	Effient	Plavix																																								
Pradaxa	Ticlid	Xaralto	Lovenox																																								
Jantoven																																											
Advil	Alka-seltzer	Anaprox	Ibuprofen																																								
Arthrotec	Vitamin E	Iron pills	Mobic																																								
Escorting	Feldene	Relafen	Multivitamins																																								
Motrin	Narcosis	Orudis	Unpin																																								
Aleve	Daypro	Iodine	Naproxen																																								
Fish oil (Omega -3)		Lodine	Piroxicam																																								
Celebrex	Maintenance medications																																										
Methotrexate	Plaquenil																																										
Seizure medications	Tylenol																																										
<i>3 DAYS PRIOR</i>	<i>2 DAYS PRIOR</i>																																										
<p>DO NOT EAT:</p> <ul style="list-style-type: none"> • cucumbers • popcorn • seeds or peels (tomatoes, grapes, seeds in breads) • nuts • corn <p>*if you want apples, pears, or peaches, you must peel them first.</p>	<ul style="list-style-type: none"> • Drink at least 2 quarts of allowed clear liquids throughout the day **NO SOLID FOOD • At 4- 6 pm, take 2 Dulcolax tablets, mix entire 238 gram bottle of Miralax prep into one 64 ounce container or two 32 ounce containers of clear, non-carbonated liquid. Drink this over a 3 hour period. 																																										
<i>1 DAY PRIOR</i>																																											
<p>CONTINUE BOWEL PREP:</p> <ul style="list-style-type: none"> • Drink at least 2 quarts of allowed clear liquids throughout the day **NO SOLID FOOD • At 4 pm, take 2 Dulcolax tablets, mix entire 238 gram bottle of Miralax prep into one 64 ounce container or two 32 ounce containers of clear, non-carbonated liquid. Drink this over a 3 hour period. 																																											

FOOD AND DRINK

ALLOWED: Clear Liquids Only. <i>We encourage you to drink plenty of the allowed liquids.</i>	NOT ALLOWED
CLEAR fruit juices without pulp : apple, white cranberry, white grape, lemonade (sweeteners are permitted)	Solid Food
Water, clear broth and bouillon (any flavor)	Alcohol
Coffee or tea (without milk or nondairy creamers)	Milk or Milk products
Gatorade, Kool-Aid, Jell-O, or other fruit flavored drinks (<i>avoid red, purple or blue food color</i>)	Orange Juice
Ice Popsicles (<i>avoid red, purple or blue food color</i>), Italian Ice	

PROCEDURE DAY

Drink only allowed clear liquids.
2 HOURS PRIOR TO ARRIVAL:
· Nothing by mouth · No gum or candy · No smoking · No water
<u>Take ALL medications as prescribed by your physician for:</u>
· cardiac · respiratory
· anti-seizure · anti-hypertensive (blood pressure)
· anti-anxiety · gastrointestinal medications
*take inhalers as prescribed and bring them with you to your appointment.
*do not take diuretic (water pills) medications

TRANSPORTATION

You MUST make arrangements for someone to drive you home, if you will be receiving any type of anesthesia or sedation. This person must be with you at check-in to confirm time for pick up. If you need to take a taxi, you MUST have a responsible person to accompany you in the taxi. **If you do not have a ride, your procedure will be cancelled.** If you are responsible for small children, make arrangements for their care during your procedure as well as for the remainder of the day.

BILLING

Please note that you will be billed separately from Amoskeag Anesthesia, PLLC for the anesthesia services provided during your procedure. Questions regarding these charges can be directed to (800) 242-1311.

INSURANCE INFORMATION

Please know your insurance guidelines! If your insurance requires a PCP authorization (insurance referral), please be sure our office has received this prior to your procedure. Please do not assume that your primary care physician's office has taken care of this. This is not necessary if your primary care physician is a Dartmouth-Hitchcock provider. Please note that we do not accept the following Medicare plans: Medicare Humana, Martin's Point Generation and New England Community Care. **Ask your insurance company:** *Am I 100% covered for a colonoscopy? Have I met my deductible and what can I expect for out of pocket expenses?*